


1.3 Physical Foundations

Living cells and organisms must perform work to stay alive and to reproduce themselves

Living Organisms Exist in a Dynamic Steady State, Never at Equilibrium with Their Surroundings



When $r_1 = r_2$
 $[\text{hemoglobin}] = \text{const}$

在動力學上。稱為 Steady State，但非 Equilibrium !

Organisms Transform Energy and Matter from Their Surroundings

Universe $\left\{ \begin{array}{l} \text{System} \\ \text{Surroundings} \end{array} \right.$

closed system : the system exchanges energy but not matter

open system : exchanges energy and matter


isolated system : exchanges neither energy nor matter

Living organism is an open system

• Living organisms create and maintain their **complex, orderly structure** using energy extracted from fuels or sunlight.

• In any physical or chemical change, the total amount of **energy in the universe remains const.**

The Flow of Electrons Provides Energy for Organisms




Energy Coupling Links Reactions in Biology

$$\Delta G = \Delta H - T\Delta S$$





Free energy (G) & Free energy change (ΔG)

Ergonic : $\Delta G < 0$ **Endergonic** : $\Delta G > 0$


(a) Mechanical example



Energy Coupling Links Reactions in Biology



a reaction :



未達平衡時 :

$$\Delta G = \Delta G^0 + RT \ln \frac{[C]^c [D]^d}{[A]^a [B]^b}$$

平衡時 : $\Delta G = 0$

$$\Delta G^0 = -RT \ln K_{eq}$$

Enzymes Promote Sequences of Chemical Reactions

Enzymes catalyze reactions by lowering the activation energy (ΔG^\ddagger)

collision frequency
 $k = zpe^{-\frac{\Delta G^\ddagger}{RT}}$
 ↓
steric factor rate const.

$e^{-\frac{\Delta G^\ddagger}{RT}}$: **fraction of collisions with sufficient energy to produce a reaction**

Each enzyme protein catalyzes a specific reaction, and each reaction in a cell is catalyzed by a different enzyme

chap 1.3 to 1.5 7

- multiplicity
- specificity
- susceptibility to regulation

↓
 give cells the capacity to lower activation barrier selectively

↓
 Effective regulation of cellular processes

↓

reaction pathway 每一步驟皆有專司負責的enzyme

Feedback inhibition

chap 1.3 to 1.5

Metabolism: Catabolism and Anabolism

<p>Nutrients Foods Solar photons</p>	<p>Cellular works Complex biomolecules Mechanical works Osmotic work</p>
--	--

catabolism (exergonic) ↓ ADP / ATP ↑

anabolism (endergonic) ↑ ADP / ATP ↓

CO₂, NH₃, H₂O precursors

chap 1.3 to 1.5 9 **Summary of 1.3**

1.4 Genetic Foundations

遺傳物質：DNA (deoxyribonucleic acid)

A. Genetic continuity is vested in DNA molecule

文明的記錄 ↓ 石頭上的文字

遺傳的記錄 ↓ DNA

reproduce themselves with nearly perfect fidelity for countless generations

chap 1.3 to 1.5 10

B. The structure of DNA allows for its repair and replication with near-perfect fidelity

雙股DNA(double helix !)

- Complementarity between the two strands accounts for the accurate replication essentials for genetic continuity
- Genetic information is encoded in the linear sequence of four kinds of subunits of DNA
- The double-helical DNA molecule contains an internal template for its own replication and repair

chap 1.3 to 1.5

The linear sequence in DNA encodes proteins with 3-dimensional structures

DNA ↓ Transcription of DNA sequence into RNA sequence

RNA ↓ Translation of RNA sequence into protein sequence

Linear amino acid sequence ↓ Noncovalent interactions

Protein with native conformation


The structure and function of proteins are encoded in its **linear a.a. sequence**

chap 1.3 to 1.5 **Summary of 1.4**


Formation of supramolecular complex

1.5 Evolutionary Foundations


Changes in the Hereditary Instructions Allow Evolution




Gene duplication and mutation (I)



Gene duplication and mutation (II)



Biomolecules First Arose by Chemical Evolution



Chemical Evolution Can Be Simulated in the Laboratory

In 1953, Stanley Muller's expt'l:

NH_3 , CH_4 , H_2 , and H_2O


Electrical sparks for a week or more

analyzed the contents:

amino acids
hydroxy acids
aldehydes
hydrogen cyanide

produced

Even, polypeptides and RNA-like molecules!! (later expt'l)



In conclusion,
many biomolecules, under prebiotic condition can be formed : including polypeptides and RNA-like molecules (as catalyst)

↓
Protein

Whether life also arose on the planets of other solar systems?




RNA or Related Precursors May Have Been the First Genes and Catalysts

RNA can act as catalysts on their own formation

↓
RNA may have been the first catalyst and first gene

chap 1.3 to 1.5

18



Comparison of prokaryotic and eukaryotic cells

Characteristic	Prokaryotic cell	Eukaryotic cell
Size	Generally small (1-10 μm)	Generally large (5-100 μm)
Genome	DNA with nonhistone protein; genome in nucleoid, not surrounded by membrane	DNA complexed with histone and nonhistone proteins in chromosomes; chromosomes in nucleus with membranous envelope
Cell division	Fission or budding; no mitosis	Mitosis, including mitotic spindle; centrioles in many species
Membrane-bounded organelles	Absent	Mitochondria, chloroplasts (in plants, some algae), endoplasmic reticulum, Golgi complexes, lysosomes (in animals), etc.
Nutrition	Absorption; some photosynthesis	Absorption, ingestion; photosynthesis in some species
Energy metabolism	No mitochondria; oxidative enzymes bound to plasma membrane; great variation in metabolic pattern	Oxidative enzymes packaged in mitochondria; more unified pattern of oxidative metabolism
Cytoskeleton	None	Complex with microtubules, intermediate filaments, actin filaments
Intracellular movement	None	Cytoplasmic streaming, endocytosis, phagocytosis, mitosis, vesicle transport

Molecular anatomy reveals evolutionary relationships

- Relatedness of species:
 - 18 century, anatomic similarities and differences among organisms (Linnaeus)
 - 19 century, phylogeny of modern organisms (Darwin)
 - 20 century, 'molecular anatomy': sequences and three-dimensional structure of nucleic acids and proteins
- Genome (the complete endowment of an organism)
- Some organisms whose genomes have been completely sequenced: (see table 1.2)

chap 1.3 to 1.5 23

TABLE 1-2 A Few of the Many Organisms Whose Genomes Have Been Completely Sequenced


Organism	Genome size (millions of nucleotide pairs)	Number of genes	Biological interest
<i>Mycoplasma genitalium</i>	0.58	483	Smallest true organism
<i>Treponema pallidum</i>	1.1	1,039	Causes syphilis
<i>Borrelia burgdorferi</i>	1.44	1,738	Causes Lyme disease
<i>Helicobacter pylori</i>	1.7	1,589	Causes gastric ulcers
<i>Methanococcus jannaschii</i>	1.7	1,783	Archaeum; grows at 85 °C
<i>Haemophilus influenzae</i>	1.8	1,738	Causes bacterial influenza
<i>Archaeoglobus fulgidus*</i>	2.2	—	High-temperature methanogen
<i>Synechocystis</i> sp.	3.6	4,003	Cyanobacterium
<i>Bacillus subtilis</i>	4.2	4,779	Common soil bacterium
<i>Escherichia coli</i>	4.6	4,377	Some strains cause toxic shock syndrome
<i>Saccharomyces cerevisiae</i>	12.5	5,770	Unicellular eukaryote
<i>Plasmodium falciparum</i>	23	5,268	Causes human malaria
<i>Caenorhabditis elegans</i>	100	19,400	Multicellular roundworm
<i>Anopheles gambiae</i>	278	13,700	Malaria vector
<i>Arabidopsis thaliana</i>	157	25,500	Model plant
<i>Oryza sativa</i>	390	37,500	Rice
<i>Drosophila melanogaster</i>	140	13,000	Laboratory fly ("fruit fly")
<i>Mus musculus domesticus</i>	2.4 × 10 ³	25,000	Laboratory mouse
<i>Pan troglodytes</i>	2.4 × 10 ³	25,000	Chimpanzee
<i>Homo sapiens</i>	2.9 × 10 ³	25,000	Human

*The number of genes is not yet determined.

Table 1-2
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company 1

Molecular phylogeny is derived from gene sequences

- When two genes DNA or protein share detectable sequence similarities, their sequences are **‘homologous’** and the proteins they encode are **‘homologs’**.
- Two homologous genes occur in the same species, they are **‘paralogous’** and the proteins are **‘paralogs’**.
- Two homologous genes occur in the different species, they are **‘orthologous’** and the proteins are **‘orthologs’**. **Annotated genome** includes (1) DNA sequence, and (2) a description of the **likely function** of each gene product (deduced from comparisons with other genomic sequences and established protein function).



Molecular phylogeny is derived from gene sequences

- The sequence differences between two homologous genes may be taken as a measure of the degree to which the two species have diverged during evolution.
- The larger the number of sequence differences, the earlier the divergence in evolutionary history.
- Phylogeny (family tree) (see fig. 1-4)